

## Remodulin - (1, 2.5, 5, 10 mg/ml INJECTABLE;IV (INFUSION), SUBC)

<b>Generic Name</b>	Treprostinil Sodium	<b>Innovator</b>	United Therapeutic Corporation
<b>Dosage</b>	1, 2.5, 5, 10 mg/ml INJECTABLE;IV (INFUSION), SUBC	<b>Branded US Sales</b>	Less Than \$1000 mn
<b>Probable FTF</b>	Less Than 5	<b>Known Para IV Filers</b>	Less Than 5
<b>Other ANDA developers</b>	More Than 5	<b>Tentative Approvals</b>	None
<b>Final Approvals</b>	Less Than 5	<b>Generic Launches</b>	Less Than 5
<b>Indication</b>	indicated for the treatment of pulmonary arterial hypertension		
<b>Complexities</b>	Yes		

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Executive Summary

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Patent Status

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Launch Timelines and Competition

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

## Remodulin - (20MG/20ML, 50MG/20ML, 100MG/20ML, 200MG/20ML)

<b>Generic Name</b>	Treprostinil Sodium	<b>Innovator</b>	None
<b>Dosage</b>	20MG/20ML, 50MG/20ML, 100MG/20ML, 200MG/20ML	<b>Branded US Sales</b>	Less Than \$1000 mn
<b>Probable FTF</b>	None	<b>Known Para IV Filers</b>	None
<b>Other ANDA developers</b>	None	<b>Tentative Approvals</b>	None
<b>Final Approvals</b>	None	<b>Generic Launches</b>	None
<b>Indication</b>	Remodulin is a prostacyclin vasodilator indicated for: • Treatment of pulmonary arterial hypertension (PAH; WHO Group 1) to diminish symptoms associated with exercise. Studies establishing effectiveness included patients with NYHA Functional Class II-IV symptoms and etiologies of idiopathic or heritable PAH (58%), PAH associated with congenital systemic-to-pulmonary shunts (23%), or PAH associated with connective tissue diseases (19%). • Patients who require transition from epoprostenol, to reduce the rate of clinical deterioration. The risks and benefits of each drug should be carefully considered prior to transition.		
<b>Complexities</b>	Yes		

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Executive Summary

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Patent Status

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Launch Timelines and Competition

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.