

Iclusig - (15, 45 mg; Tablet)

| | | | |
|------------------------------|---|-----------------------------|---------------------|
| Generic Name | Ponatinib | Innovator | Takeda |
| Dosage | 15, 45 mg; Tablet | Branded US Sales | Less Than \$1000 mn |
| Probable FTF | Less Than 5 | Known Para IV Filers | Less Than 5 |
| Other ANDA developers | Less Than 5 | Tentative Approvals | Less Than 5 |
| Final Approvals | Less Than 5 | Generic Launches | None |
| Indication | Iclusig is a kinase inhibitor indicated for the treatment of adult patients with chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ALL) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy (1). This indication is based upon response rate. There are no trials verifying an improvement in disease-related symptoms or increased survival with Iclusig. | | |
| Complexities | Yes | | |

Chronology Of Events

Please Contact contact@researchdelta.com to get Detailed Information.

Executive Summary

Please Contact contact@researchdelta.com to get Detailed Information.

Patent Status

Please Contact contact@researchdelta.com to get Detailed Information.

Launch Timelines and Competition

Please Contact contact@researchdelta.com to get Detailed Information.

Chronology Of Events

Please Contact contact@researchdelta.com to get Detailed Information.

Iclusig - (10, 30 mg; Tablet)

| | | | |
|------------------------------|--|-----------------------------|---------------------|
| Generic Name | Ponatinib | Innovator | Takeda |
| Dosage | 10, 30 mg; Tablet | Branded US Sales | Less Than \$1000 mn |
| Probable FTF | Less Than 5 | Known Para IV Filers | Less Than 5 |
| Other ANDA developers | Less Than 5 | Tentative Approvals | Less Than 5 |
| Final Approvals | None | Generic Launches | None |
| Indication | Iclusig is a kinase inhibitor indicated for the treatment of adult patients with chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ALL) that is resistant or intolerant to prior tyrosine kinase inhibitor therapy | | |
| Complexities | Yes | | |

Chronology Of Events

Please Contact contact@researchdelta.com to get Detailed Information.

Executive Summary

Please Contact contact@researchdelta.com to get Detailed Information.

Patent Status

Please Contact contact@researchdelta.com to get Detailed Information.

Launch Timelines and Competition

Please Contact contact@researchdelta.com to get Detailed Information.

Chronology Of Events

Please Contact contact@researchdelta.com to get Detailed Information.