

## Xeljanz - (5 mg; Tablet, Oral)

<b>Generic Name</b>	Tofacitinib	<b>Innovator</b>	Pfizer
<b>Dosage</b>	5 mg; Tablet, Oral	<b>Branded US Sales</b>	Less Than \$1000 mn
<b>Probable FTF</b>	Less Than 5	<b>Known Para IV Filers</b>	More Than 5
<b>Other ANDA developers</b>	More Than 5	<b>Tentative Approvals</b>	More Than 5
<b>Final Approvals</b>	Less Than 5	<b>Generic Launches</b>	None
<b>Indication</b>	Used to treat moderate to severe rheumatoid arthritis or active psoriatic arthritis (PsA) in adults who have tried methotrexate other medications without successful treatment of symptoms		
<b>Complexities</b>	Yes		

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Executive Summary

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Patent Status

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Launch Timelines and Competition

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

## Xeljanz - (10 mg; Tablet, Oral)

<b>Generic Name</b>	Tofacitinib	<b>Innovator</b>	Pfizer
<b>Dosage</b>	10 mg; Tablet, Oral	<b>Branded US Sales</b>	Less Than \$1000 mn
<b>Probable FTF</b>	Less Than 5	<b>Known Para IV Filers</b>	More Than 5
<b>Other ANDA developers</b>	More Than 5	<b>Tentative Approvals</b>	Less Than 5
<b>Final Approvals</b>	None	<b>Generic Launches</b>	None
<b>Indication</b>	Xeljanz is indicated for the treatment of adult patients with active psoriatic arthritis (PsA) who have had an inadequate response or intolerance to methotrexate or other disease-modifying antirheumatic drugs.		
<b>Complexities</b>	Yes		

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Executive Summary

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Patent Status

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Launch Timelines and Competition

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

## Xeljanz - (1 mg/mL ; Oral Solution)

<b>Generic Name</b>	Tofacitinib	<b>Innovator</b>	Pfizer
<b>Dosage</b>	1 mg/mL ; Oral Solution	<b>Branded US Sales</b>	Less Than \$1000 mn
<b>Probable FTF</b>	Less Than 5	<b>Known Para IV Filers</b>	Less Than 5
<b>Other ANDA developers</b>	Less Than 5	<b>Tentative Approvals</b>	Less Than 5
<b>Final Approvals</b>	Less Than 5	<b>Generic Launches</b>	None
<b>Indication</b>	<p>XELJANZ/XELJANZ XR/XELJANZ Oral Solution is a Janus kinase (JAK) inhibitor indicated for: 1) Rheumatoid Arthritis - XELJANZ/XELJANZ XR is indicated for the treatment of adult patients with moderately to severely active rheumatoid arthritis who have had an inadequate response or intolerance to methotrexate. 2) Psoriatic Arthritis - XELJANZ/XELJANZ XR is indicated for the treatment of adult patients with active psoriatic arthritis who have had an inadequate response or intolerance to methotrexate or other disease-modifying antirheumatic drugs (DMARDs). ? Limitations of Use: Use of XELJANZ/XELJANZ XR in combination with biologic DMARDs or potent immunosuppressants such as azathioprine and cyclosporine is not recommended. 3) Ulcerative Colitis - XELJANZ/XELJANZ XR is indicated for the treatment of adult patients with moderately to severely active ulcerative colitis (UC), who have had an inadequate response or who are intolerant to TNF blockers. 4) Polyarticular Course Juvenile Idiopathic Arthritis - XELJANZ/XELJANZ Oral Solution is indicated for the treatment of active polyarticular course juvenile idiopathic arthritis (pcJIA) in patients 2 years of age and older.</p>		
<b>Complexities</b>	Yes		

### Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Executive Summary

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Patent Status

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

### Launch Timelines and Competition

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.

## Chronology Of Events

Please Contact [contact@researchdelta.com](mailto:contact@researchdelta.com) to get Detailed Information.